arXiv:1202.6261v1 [math.AG] 28 Feb 2012

GLOBALLY GENERATED VECTOR BUNDLES ON $\mathbb{P}^n$ WITH $c_1 = 3$

CRISTIAN ANGHEL AND NICOLAE MANOLACHE

Abstract. One classifies the globally generated vector bundles on $\mathbb{P}^n$ whith the first Chern class $c_1 = 3$ (in this paper $n \neq 3$, the case $n = 3$ being studied in [18]). The case $c_1 = 1$ is very easy, the case $c_1 = 2$ was done in [28], the case $c_1 = 3$, rank= 2 was settled in [13] and the case $c_1 \leq 5$, rank=2 in [6]. Our work is based on Serre's theorem relating vector bundles of rank = 2 with codimension 2 lci subschemes and its generalization for higher ranks, considered firstly by Vogelaar in [33].

1. Introduction

We use freely the standard notation, namely that introduced in [25], used (and extended in [7]) (cf. also [9]). Our base field is $\mathbb{C}$.

A globally generated nontrivial vector bundle E on a a projective space has the first Chern class $c_1(E) > 0$. When $c_1(E) = 1$, modulo a trivial summand, $E \cong \mathcal{O}(1)$ or $E \cong T(-1)$, T being the tangent bundle. The case $c_1(E) = 2$ is studied in [28] and the case $c_1 = 3$ on $\mathbb{P}^3$ was studied in [18]. Globally generated rank 2 vector bundles with $c_1 = 3$ are listed in [13]. Rank 2 globally generated vector bundles on $\mathbb{P}^n$, $n \geq 3$, with $c_1 \leq 5$ are listed in [6].

The aim of this paper is to prove:

Theorem. *(i) On $\mathbb{P}^2$ the globally generated vector bundles of rank 2 with the first Chern class $c_1 = 3$, which are not direct sums of line*

bundles are as follows:

$$0 \to \mathcal{O}(-1) \to 2\mathcal{O} \oplus \mathcal{O}(2) \to E \to 0 \tag{1}$$

$$E \cong \Omega^1(3) \cong T_{\mathbb{P}^2} \tag{2}$$

$$0 \to \mathcal{O}(-1) \to \mathcal{O} \oplus 2\mathcal{O}(1) \to E \to 0 \tag{3}$$

$$0 \to 2\mathcal{O}(-1) \to 3\mathcal{O} \oplus \mathcal{O}(1) \to E \to 0 \tag{4}$$

$$0 \to \mathcal{O}(-2) \to 2\mathcal{O} \oplus \mathcal{O}(1) \to E \to 0 \tag{5}$$

$$0 \to \mathcal{O}(-3) \to 3\mathcal{O} \to E \to 0 \tag{6}$$

$$0 \to \mathcal{O}(-2) \oplus \mathcal{O}(-1) \to 4\mathcal{O} \to E \to 0 \tag{7}$$

$$0 \to 3\mathcal{O}(-1) \to 5\mathcal{O} \to E \to 0 \tag{8}$$

(ii) On $\mathbb{P}^4$ *the globally generated vector bundles with* $c_1 = 3$ *of rank at most* 4 *are:*

– *none of rank* 2

– *of rank* 3*: the dual of the Trautmann-Vetter (Tango) bundle:*

$$0 \to T(-2) \to 7\mathcal{O} \to E \to 0$$

– *of rank* 4*: given by exact sequences:*

$$0 \to \mathcal{O}(-1) \to 4\mathcal{O} \oplus \mathcal{O}(2) \to E \to 0 \tag{1}$$

$$0 \to \mathcal{O}(-1) \to 3\mathcal{O} \oplus 2\mathcal{O}(1) \to E \to 0 \tag{2}$$

$$0 \to 2\mathcal{O}(-1) \to 5\mathcal{O} \oplus \mathcal{O}(1) \to E \to 0 \tag{3}$$

$$0 \to \mathcal{O}(-2) \to 4\mathcal{O} \oplus \mathcal{O}(1) \to E \to 0 \tag{4}$$

$$0 \to 3\mathcal{O}(-1) \to 7\mathcal{O} \to E \to 0 \tag{5}$$

$$0 \to \mathcal{O}(-2) \oplus \mathcal{O}(-1) \to 6\mathcal{O} \to E \to 0 \tag{6}$$

$$0 \to \mathcal{O}(-3) \to 5\mathcal{O} \to E \to 0 \tag{7}$$

$$E \cong \Omega^1(2) \tag{8}$$

(iii) On $\mathbb{P}^5$ *the globally generated vector bundles with* $c_1 = 3$ *of rank at most* 5 *are:*

– *none of ranks* 2, 3, 4

– *of rank* 5 *given by exact sequences:*

$$0 \to \mathcal{O}(-1) \to 5\mathcal{O} \oplus \mathcal{O}(2) \to E \to 0 \tag{1}$$
$$0 \to \mathcal{O}(-1) \to 4\mathcal{O} \oplus 2\mathcal{O}(1) \to E \to 0 \tag{2}$$
$$0 \to 2\mathcal{O}(-1) \to 6\mathcal{O} \oplus \mathcal{O}(1) \to E \to 0 \tag{3}$$
$$0 \to \mathcal{O}(-2) \to 5\mathcal{O} \oplus \mathcal{O}(1) \to E \to 0 \tag{4}$$
$$0 \to 3\mathcal{O}(-1) \to 8\mathcal{O} \to E \to 0 \tag{5}$$
$$0 \to \mathcal{O}(-2) \oplus \mathcal{O}(-1) \to 7\mathcal{O} \to E \to 0 \tag{6}$$
$$0 \to \mathcal{O}(-3) \to 6\mathcal{O} \to E \to 0 \tag{7}$$

(iv) On $\mathbb{P}^n$, $n = 2, 4, 5$ *the globally generated vector bundles of rank* $> n$ *with* $c_1 = 3$ *are extensions of those of rank* n *by trivial bundles.*

(v) On $\mathbb{P}^n$, $n \geq 5$ *the globally generated vector bundles with* $c_1 = 3$ *have rank* $r \geq n$ *and are given by exact sequences:*

$$0 \to \mathcal{O}(-1) \to r\mathcal{O} \oplus \mathcal{O}(2) \to E \to 0 \tag{1}$$
$$0 \to \mathcal{O}(-1) \to (r-1)\mathcal{O} \oplus 2\mathcal{O}(1) \to E \to 0 \tag{2}$$
$$0 \to 2\mathcal{O}(-1) \to (r+1)\mathcal{O} \oplus \mathcal{O}(1) \to E \to 0 \tag{3}$$
$$0 \to \mathcal{O}(-2) \to r\mathcal{O} \oplus \mathcal{O}(1) \to E \to 0 \tag{4}$$
$$0 \to 3\mathcal{O}(-1) \to (r+3)\mathcal{O} \to E \to 0 \tag{5}$$
$$0 \to \mathcal{O}(-2) \oplus \mathcal{O}(-1) \to (r+2)\mathcal{O} \to E \to 0 \tag{6}$$
$$0 \to \mathcal{O}(-3) \to (r+1)\mathcal{O} \to E \to 0 \tag{7}$$

Aknowledgements. *The starting point of this paper was a discussion with Iustin Coandă about* [28]. *We want to thank him for this, for his interest, for pointing us some useful references and also for useful comments on a previous version of this paper. The second author expresses his thanks to the Institute of Mathematics, Oldenburg University, especially to Udo Vetter, for warm hospitality. In writing this paper the authors had no other support besides the membership to the Institute of Mathematics "Simion Stolow" of the Romanian Academy.*

2. Preliminaries

We recall some general results, mostly of which are the starting remarks in [28]:

A. (J.-P. Serre, cf. [2]) If a vector bundle E or rank $r > n$ on a smooth variety X of dimension n is globally generated, then E contains a trivial subbundle of rank $= r - n$.

B. If a globally generated vector bundle E or rank $r \leq n$ on $\mathbb{P}^n$ has $c_r = 0$, then E contains a trivial subbundle of rank $= 1$ (cf. [22], chap. I, Lemma 4.3.2).

C. If E is a globally generated vector bundle on $\mathbb{P}^n$, then E contains a trivial direct summand of rank $= \dim H^0(E^\vee)$ (cf. [23], Lemma 3.9).

D. If E is a vector bundle on $\mathbb{P}^n$, which is globally generated and for which $H^0(E(-c_1)) \neq 0$, where c_1 is the first Chern class of E, then, modulo a trivial summand, $E \cong \mathcal{O}(c_1)$ (cf. [27], Prop.1).

E. (Vogelaar, Arrondo's setting, cf. [1], Theorem 1.1)

(i) If E is a vector bundle of rank r on $\mathbb{P}^n$ and $r-1$ sections $s_1, \ldots, s_{r-1}$ of E has as dependency locus a subscheme Y of $\mathbb{P}^n$ which is lci, then one has an exact sequence:

$$0 \to (r-1)\mathcal{O} \xrightarrow{(s_1,\ldots,s_{r-1})} E \to I_Y \otimes L \to 0 \quad ,$$

where $L = \Lambda^r E$. In this case, if N is the normal sheaf of Y, then $\Lambda^2 N \otimes L^\vee|_Y$ is generated by $r-1$ global sections.

(ii) Conversely, if Y is a lci subscheme of codimension 2 of $\mathbb{P}^n$ and L is a line bundle on $\mathbb{P}^n$ with the property $H^2(L^\vee) = 0$, if $\Lambda^2 N \otimes L^\vee|_Y$ is generated by $r-1$ global sections, then there exists a vector bundle E of rank r such that $\Lambda^r E = L$ and E has $r-1$ global sections whose dependence locus is Y.

F. If $\varphi : F \to G$ is a general morphism between vector bundles over a smooth variety and $\mathcal{H}om(F, G)$ is globally generated, then the dependency locus Y of φ is either empty or has codimension $\operatorname{rank}(G) - \operatorname{rank}(F) + 1$ and is nonsingular outside a subset of codimension $\geq \operatorname{rank}(G) - \operatorname{rank}(F) + 3$ (cf. [3] 4.1 or [5] for a more general result and [24], [26] for the case of arithmetically Cohen-Macaulay curves in $\mathbb{P}^3$).

□

According to **A**, any vector bundle on $\mathbb{P}^n$ of rank $> n$ is an extension of a rank n vector bundle by a trivial one, so on $\mathbb{P}^n$, modulo extensions by trivial bundles, it is enough to find the globally generated vector bundles of rank at most n with $c_1 = 3$.

3. Vector Bundles on $\mathbb{P}^2$

In this case we have to treat only the case of rank =2. This is treated in [13] and we give here other arguments. Take E a globally generated vector bundle of rank 2 on $\mathbb{P}^2$, non-splittable in line bundles, with $c_1(E) = 3$; we may assume $H^0(E(-3)) = 0$, according to **D**. If no section of it has an empty zero's set (in which case E splits), a general section of it has a smooth zero set Z which according to [12], Prop. 5.1, or [10], Prop. 1.4. is a union of reduced closed points. One has an exact sequence:

$$(*) \qquad 0 \to \mathcal{O} \to E \to I_Z(3) \to 0$$

I. **Assume** $H^0(E(-3)) = 0$, $H^0(E(-2)) \neq 0$. The exact sequence (*) shows that Z lies on a line. As $I_Z(3)$ is generated by global sections, Z is included in a complete intersection of type $(1, 3)$, so $c_2(E) = \deg(Z) \leq 3$. On the other hand, as E does not split, one has an exact sequence:

$$0 \to \mathcal{O} \to E(-2) \to I_X(-1) \to 0 \quad ,$$

where X is lci subscheme of dimension 0 of $\mathbb{P}^2$. One has $c_2(E(-2)) = c_2(E) - 2 = \deg(X) \geq 1$. It follows $c_2(E) = 3$ so that Z is a complete intersection of type $(1, 3)$ and one gets for E a resolution of the shape:

$$0 \to \mathcal{O}(-1) \to 2\mathcal{O} \oplus \mathcal{O}(2) \to E \to 0 \quad .$$

II. **Assume** $H^0(E(-2)) = 0$, $H^0(E(-1)) \neq 0$. In this case E is a stable vector bundle of rank 2 on $\mathbb{P}^2$. Let $F = E(-2)$. Then $c_1(F) = -1$ and, because of stability, $c_2(F) \geq 1$. If $c_2(F) = n$, then F is given by a monad (cf. [14]):

$$(n-1)\mathcal{O}(-1) \xrightarrow{a} n\Omega(1) \xrightarrow{a^T(-1)} (n-1)\mathcal{O} \quad .$$

When $n = 1$, $E \cong \Omega(3) \cong T_{\mathbb{P}^2}$. From now on $n \geq 2$.

As $c_1(E(-1)) = 1$, we have an exact sequence:

$$(**) \qquad 0 \to \mathcal{O} \to E(-1) \to I_X(1) \to 0 \quad ,$$

where X is a subscheme of dimension 0 of $\deg(X) = c_2(E(-1)) = n$.

In the exact sequence (*) the scheme Z consists of $c_2(E) = n + 2$ points. The condition that Z lies in a complete intersection of type $(2, 3)$ gives $n + 2 \leq 6$, so $2 \leq n \leq 4$.

If $n = 2$, then Z is a complete intersection of type $(2, 2)$, and one gets an exact sequence:

$$0 \to \mathcal{O}(-1) \to \mathcal{O} \oplus 2\mathcal{O}(1) \to E \to 0 \quad .$$

If n=3 then X cannot lie on a line. Indeed, $h^0(I_X(1) \neq 0$ would give, via (**), X being a complete intersection of type $(1, 3)$, $h^0(E(-1)) = 2$,

$h^0(E) = 6$. From(*) would follow that Z, a set of 5 different points, would lie on two distinct conics, contradiction. It follows that X has a resolution of the shape:

$$0 \to 2\mathcal{O}(-3) \to 3\mathcal{O}(-2) \to I_X \to 0 \ ,$$

where from E has a resolution:

$$0 \to 2\mathcal{O}(-1) \to 3\mathcal{O} \oplus \mathcal{O}(1) \to E \to 0$$

Remark. The same result could be obtained using the resolution of I_Z, Z being a set of 5 points not situated on a line:

$$0 \to 2\mathcal{O}(-4) \to 2\mathcal{O}(-3) \oplus \mathcal{O}(-2) \to I_Z \to 0 \ .$$

If $n = 4$, then Z is a complete intersection of type $(2, 3)$ and for E follows a resolution of the shape:

$$0 \to \mathcal{O}(-2) \to 2\mathcal{O} \oplus \mathcal{O}(1) \to E \to 0 \ .$$

III. **Assume** $H^0(E(-1)) = 0$. With the notation from above, if $c_1(F) = 1$, $c_2(F) = n \geq 2$ then $\deg(Z) = n + 2 \leq 9$, Z being included in a complete intersection of type $(3, 3)$, and so $n \leq 7$. As Z lies on no conic, $4 \leq n \leq 7$.

For $n = 4, 5, 6$ i.e. $\deg(Z) = 6, 7, 8$ and the resolutions of I_Z are:

(a) $deg(Z) = 6$:

$$0 \to 3\mathcal{O}(-4) \to 4\mathcal{O}(-3) \to I_Z \to 0$$

and then the resolution of E is:

$$0 \to 3\mathcal{O}(-1) \to 5\mathcal{O} \to E \to 0 \ .$$

(b) $\deg(Z) = 7$:

$$0 \to \mathcal{O}(-5) \oplus \mathcal{O}(-4) \to 3\mathcal{O}(-3) \to I_Z \to 0$$

and then the resolution of E is:

$$0 \to \mathcal{O}(-2) \oplus \mathcal{O}(-1) \to 4\mathcal{O} \to E \to 0 \ ;$$

(c) $deg(Z) = 8$:

$$0 \to 2\mathcal{O}(-5) \to \mathcal{O}(-4) \oplus 2\mathcal{O}(-3) \to I_Z \to 0$$

and then the resolution of E is:

$$0 \to 2\mathcal{O}(-2) \to \mathcal{O}(-1) \oplus 3\mathcal{O} \to E \to 0 \ .$$

In the last case E is not globally generated.

To prove (a), use the fact that Z is linked in a complete intersection of type $(3, 3)$ to a scheme of degree 3. In general this can be: three non-collinear simple points, three collinear simple points (here are included the cases when two or all three coincide), a double point and a simple one but not on a line, the first infinitesimal neighbourhood of a point.

Applying [24], 4.1, one gets the resolution of I_Z from the resolution of the linked scheme. In our case, when the linked scheme is contained in a line, Z would be contained in a conic, what we excluded. In the other cases one obtains the asserted shape. The same result can be obtained from [8], where more complicated cases (6 fat points!) are studied.

In the same way one proves the cases (b), (c).

If $deg(Z) = 9$, then Z is a complete intersection of two cubics and the resolution of E is:

$$0 \to \mathcal{O}(-3) \to 3\mathcal{O} \to E \to 0 \quad ,$$

4. Vector Bundles on $\mathbb{P}^4$ or $\mathbb{P}^5$

We do not have indecomposable rank 2 vector bundles with $c_1 = 3$ on $\mathbb{P}^3$ (cf. [18] or [6]) so that, applying Horrocks' criterion (cf. [11], or [22], 2.3.2) they do not exist on $\mathbb{P}^n$, $n \geq 3$, either. In this section $\mathbb{P}^n$ is $\mathbb{P}^4$ or $\mathbb{P}^5$. We use **E.** and **F.** If E is a globally generated vector bundle of rank r on $\mathbb{P}^n$, with $c_1(E) = 3$ and if E has no subbundle of rank 1, then one has an exact sequence:

$$(***) \qquad 0 \to (r-1)\mathcal{O} \to E \to I_Z(3) \to 0 \quad ,$$

where Z is a smooth subvariey of $\mathbb{P}^n$. Observe $\Lambda_Z := \Lambda^2 N \otimes L^\vee | Z \cong \omega_Z \otimes \omega_{\mathbb{P}^n}^{-1} \otimes L^\vee | Z \cong \omega_Z(n-2)$. Let E be non-splittable.

I. **Assume** $H^0(E(-3)) = 0$, $H^0(E(-2)) \neq 0$. If $E(-2)$ has a section without zeros, then E is an extension of vector bundles:

$$0 \to \mathcal{O}(2) \to E \to E' \to 0$$

where E' is globally generated and has $c_1 = 1$. Then, modulo a trivial summand, E' is $\mathcal{O}(1)$ or $T(-1)$. In the first case E is splittable, in the second has rank $> n$. Suppose now that E has no rank 1 subbundle. According to **E.** and **F.**, one has an extension:

$$0 \to (r-1)\mathcal{O} \to E \to I_Z(3) \to 0 \quad ,$$

with Z a smooth codimension 2 subvariety of $\mathbb{P}^n$. In our hypothesis, Z is contained in a hyperplane, and I_Z is generated by cubic forms so that Z is either a linear subspace, a complete intersection of type $(1,2)$ or of type $(1,3)$. When Z is a linear subspace $\Lambda_Z \cong \mathcal{O}(-1)$ is not globally generated. When Z is a complete intersection of type $(1,2)$ $\Lambda_Z \cong \mathcal{O}_Z$. Then $E \cong \mathcal{O}(1) \oplus \mathcal{O}(2)$. When Z is a complete intersection of type $(1,3)$, $\Lambda_Z \cong \mathcal{O}(1)$ is generated by $(n-1)$ sections, so that E is given by an extension:

$$0 \to (n-1)\mathcal{O} \to E \to I_Z(3) \to 0 \quad .$$

Then E is given by an extension:

$$0 \to \mathcal{O}(-1) \to n\mathcal{O} \oplus \mathcal{O}(2) \to E \to 0 \quad .$$

II. **Assume** $H^0(E(-2)) = 0$ and $H^0(E(-1)) \neq 0$.

Then in the extension (***) Z is not contained in a hyperplane, is contained in at least one quadric and I_Z is generated by cubics. Then $3 \leq \deg(Z) \leq 6$.

(a) $\deg(Z) = 3$. According to [34], Z is one of:

(a1) $Z \subset \mathbb{P}^5$, image of a Segre embedding of $\mathbb{P}^1 \times P^2$ in P^5,

(a2) $Z \subset \mathbb{P}^4$, a smooth hyperplane section of the first case.

The case (a1) does not give an indecomposable vecor bundle. Indeed, I_Z would have a resolution of the shape:

$$0 \to 2\mathcal{O}_{\mathbb{P}^5}(-3) \to 3\mathcal{O}_{\mathbb{P}^5}(-2) \to I_Z \to 0 \quad ,$$

(cf. [4]) and $\Lambda_Z|_{\mathbb{P}^1\times\mathbb{P}^2} \cong \mathcal{O}_{\mathbb{P}^1\times\mathbb{P}^2}(1,0)$. It is easy to see that, in this case, $E \cong 3\mathcal{O}(1)$.

In the case (a2) one gets for $Z' = Z \cap \mathbb{P}^4$ a similar resolution:

$$0 \to 2\mathcal{O}_{\mathbb{P}^4}(-3) \to 3\mathcal{O}_{\mathbb{P}^4}(-2) \to I_{Z'} \to 0 \quad ,$$

and again E is direct sum of line bundles.

(b) $\deg(Z) = 4$. The only codimension 2, smooth projective varieties of degree 4, not contained in a hyperplane, contained in a hyperquadric and with the ideal generated by $H^0(I_Z(3))$ are the complete intersections of type $(2,2)$, (cf. [29], or [20]). In this case one gets a resolution for E:

$$0 \to \mathcal{O}(-1) \to (n-1)\mathcal{O} \oplus 2\mathcal{O}(1) \to E \to 0 \quad .$$

(c) $\deg(Z) = 5$. The only codimension 2, smooth projective varieties of degree 5, not contained in a hyperplane, with $H^0(I_Z(2)) \neq 0$ and with the ideal generated by degree 3 component are (cf. [20], [19] or [15]):

(c1) Castelnuovo surface $Z \cong \mathbb{P}^2(x_0, \ldots, x_7)$, i.e. P^2 blown-up in 8 points in general position, embedded in $\mathbb{P}^4$ via the quartics through x_i, with multiplicity 2 in x_0. I_Z has a minimal resolution of the shape (see [20], proof of 2.9):

$$0 \to 2\mathcal{O}(-4) \to 2\mathcal{O}(-3) \oplus \mathcal{O}(-2) \to I_Z \to 0 \quad .$$

$\Lambda_Z \cong \omega_Z(2)$ is generated by global sections. As $h^0(\omega_Z(3)) = 8$, one gets a vector bundle F of rank 9 with a resolution of the shape:

$$0 \to 2\mathcal{O}(-1) \to 10\mathcal{O} \oplus \mathcal{O}(1) \to F \to 0 \quad .$$

F is an extension by a trivial bundle of a vector bundle E of rank $= 4$. Indeed, by the property **A.**, F contains a trivial subbundle of rank 5. Let E be the quotient. By standard arguments one sees that this lifts

to a trivial subbundle of $10\mathcal{O}\oplus\mathcal{O}(1)$ such that one has the commutative diagram;

$$\begin{array}{ccccccccc}
 & & & & 0 & & 0 & & \\
 & & & & \downarrow & & \downarrow & & \\
 & & & & 5\mathcal{O} & = & 5\mathcal{O} & & \\
 & & & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & 2\mathcal{O}(-1) & \longrightarrow & 10\mathcal{O}\oplus\mathcal{O}(1) & \longrightarrow & F & \longrightarrow & 0 \\
 & & \| & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & 2\mathcal{O}(-1) & \longrightarrow & 5\mathcal{O}\oplus\mathcal{O}(1) & \longrightarrow & E & \longrightarrow & 0 \\
 & & & & \downarrow & & \downarrow & & \\
 & & & & 0 & & 0 & &
\end{array}$$

(c2) Z = Castelnuovo threefold in $\mathbb{P}^5$. In this case the resolution of I_Z has the shape (cf. [19], proof of 1.5):

$$0 \to 2\mathcal{O}(-4) \to 2\mathcal{O}(-3)\oplus\mathcal{O}(-2) \to I_Z \to 0 \quad .$$

Like above, one obtains a vector bundle E on $\mathbb{P}^5$ with a resolution of the form:

$$0 \to 2\mathcal{O}(-1) \to 6\mathcal{O}\oplus\mathcal{O}(1) \to E \to 0 \quad .$$

(d) $\deg(Z) = 6$. The condition that Z is contained in a quadric makes possible only: Z = complete intersection of type $(2,3)$. By standard arguments one obtains a vector bunlde E on $\mathbb{P}^n$ of rank $= n$ with a resolution of the shape:

$$0 \to \mathcal{O}(-2) \to n\mathcal{O}\oplus\mathcal{O}(1) \to E \to 0 \quad .$$

III. **Assume** $H^0(E(-1)) = 0$. We have to consider the smooth projective varieties of codimension 2, not contained in any hyperquadric, but with ideal generated by cubic forms.

(a) $\deg(Z) = 4$. The only possibility is the Veronese surface in $\mathbb{P}^4$. It is known (cf. [20], Corr. 2.8) that the ideal of Z has an Ω-resolution of the form:

$$0 \to 3\mathcal{O} \to \Omega^1(2) \to I_Z(3) \to 0 \quad .$$

In fact any vector bundle E in an extension:

$$0 \to 3\mathcal{O} \to E \to I_Z(3) \to 0 \quad .$$

with Z the Veronese surface in $\mathbb{P}^4$ is isomorphic to $\Omega^1(2)$. This can be seen via the Beilinson spectral sequence (cf. [22], II, 3.1.3) associated to $E(-1)$, which turns out to be (very) degenerated.

(b) $\deg(Z) = 5$. In our conditions Z is an elliptic scroll with $e = -1$ and $H \equiv C + 2f$ (cf. [20], Theorem 2.10). In [20] it is shown that I_Z has a resolution:

$$0 \to T_{\mathbb{P}^4}(-2) \to 5\mathcal{O} \to I_Z(3) \to 0$$

From here one gets a rank 3 vector bundle as an extension:

$$0 \to 2\mathcal{O} \to E \to I_Z(3) \to 0$$

The resolution of $I_Z(3)$ provides a resolution of E:

$$0 \to T_{\mathbb{P}^4}(-2) \to 7\mathcal{O} \to E \to 0 \quad ,$$

which shows that E is the dual of a Trautmann-Vetter (Tango) bundle on $\mathbb{P}^4$. (cf. [31], [32], [30], [16]).

(c) $\deg(Z) = 6$. The only possibilities are the Bordiga surface in $\mathbb{P}^4$ and the Bordiga threefold in $\mathbb{P}^5$ (cf. [19] or [15]).

In both cases the minimal resolution has the shape:

$$0 \to 3\mathcal{O}(-4) \to 4\mathcal{O}(-3) \to I_Z \to 0 \quad .$$

These two cases will produce a rank 4 bundle on $\mathbb{P}^4$ with a resolution:

$$0 \to 3\mathcal{O}_{\mathbb{P}^4}(-1) \to 7\mathcal{O}_{\mathbb{P}^4} \to E \to 0$$

and a rank 5 bundle on $\mathbb{P}^5$ with a resolution:

$$0 \to 3\mathcal{O}_{\mathbb{P}^5}(-1) \to 8\mathcal{O}_{\mathbb{P}^5} \to E \to 0 \quad .$$

(d) $\deg(Z) = 7$. We could use [21], but one can proceed directly. Indeed, in our situation Z is linked in a complete intersection of type $(3, 3)$ to a locally Cohen-Macaulay subscheme Y of $\mathbb{P}^n$, $(n = 4, 5)$, of pure codimension 2 and degree 2. Y cannot be a double structure on a linear subspace of codimension 2, as such a structure does not exist (cf. [17]) and cannot be a union of two linear subspaces of codimension 2 meeting along a linear subsapce of codimension 4, as this would contradict Cohen-Macaulay property. Then Y is a quadric of codimension 2, degenerated or not. It follows the resolution of I_Z:

$$0 \to \mathcal{O}(-5) \oplus \mathcal{O}(-4) \to 3\mathcal{O}(-3) \to I_Z \to 0$$

These Z's give rise to the vector bundles on $\mathbb{P}^4$, respectively $\mathbb{P}^5$:

$$0 \to \mathcal{O}_{\mathbb{P}^4}(-2) \oplus \mathcal{O}_{\mathbb{P}^4}(-1) \to 6\mathcal{O}_{\mathbb{P}^4} \to E \to 0 \quad ,$$

$$0 \to \mathcal{O}_{\mathbb{P}^5}(-2) \oplus \mathcal{O}_{\mathbb{P}^5}(-1) \to 7\mathcal{O}_{\mathbb{P}^5} \to E \to 0 \quad ,$$

of ranks 4, respectively 5.

(e) $\deg(Z) = 8$. This situation does not occur in our context, as such a Z, being linked to linear subspace would have $I_Z(3)$ not globally generated.

(f) $\deg(Z) = 9$. Then Z is a complete intersection and the vector bundles obtained this way are:

$$0 \to \mathcal{O}_{\mathbb{P}^4}(-3) \to 5\mathcal{O}_{\mathbb{P}^4} \to E \to 0 \quad ,$$

of rank 4 on $\mathbb{P}^4$ and

$$0 \to \mathcal{O}_{\mathbb{P}^5}(-3) \to 6\mathcal{O}_{\mathbb{P}^5} \to E \to 0 \quad ,$$

of rank 5 on $\mathbb{P}^5$.

5. Vector Bundles on $\mathbb{P}^n$, $n \geq 5$

From (iii) and (iv) in the Theorem it follows directly (v) for $\mathbb{P}^5$. We prove (v) by induction by standard (known) cohomological arguments.

Lemma 5.1. *If E is a vector bundle on $\mathbb{P}^{n+1}$ and its restriction E' to a hyperplane has $H^i(E'(j)) = 0$ for $1 \leq i \leq n-2$ and any j, then $H^i(E(j)) = 0$ for $1 \leq i \leq n-1$ and any j.*

Proof. From the exact sequence:

$$0 \to E(-1) \to E \to E' \to 0$$

it follows, for all j:

$$\begin{aligned} H^i(E(j-1)) \cong H^i(E(j)), \quad & 2 \leq i \leq n-2 \quad , \\ H^1(E(j-1)) \to H^1(E(j)) \quad & \text{surjective} \\ H^{n-1}(E(j-1)) \to H^{n-1}(E(j)) \quad & \text{injective} \end{aligned}$$

As $H^i(E(j)) = 0$ for $i = 1, \ldots n$ and $j \ll 0$ or $j \gg 0$ the assertion follows. □

Lemma 5.2. *If E is a vector bundle on $\mathbb{P}^n$ with the property $H^i(E(j)) = 0$ for $1 \leq i \leq n-2$ then it has a resolution of length 1:*

$$0 \to L_1 \to L_0 \to E \to 0 \quad ,$$

where L_0, L_1 are direct sums of line bundles.

Proof. Indeed, take a (minimal) surjection $L_0 \to E \to 0$, with L_0 a direct sum of line bundles and denote by L_1 the kernel. It follows $H^i((L_1(j)) = 0$ for $i = 1, \ldots n-1$ and all j. By the spliting criterion of Horrocks (cf. [11], or [22] I, Theorem 2.3.1) it follows that L_1 is a direct sum of line bundles. □

The assertion (v) in the Theorem follows directly from the above lemmas.

References

[1] E. Arrondo, A Home Made Hartshorne-Serre Correspondence, Rev. Mat. Complut, **20,** (2007), no. 2, 423-443

[2] M.F. Atiyah, Vector Bundles over an Elliptic Curve, Proc.London Math. Soc. **3** 7, 1957, 414-452

[3] C. Bănică, Smooth Reflexive Sheaves, Rev. Roum. Math. Pures et Appl., **36**, (1991), 571-593

[4] Ş. Bărcănescu, N. Manolache, Nombres de Betti d'une singularité de Segre-Veronese, C.R. Sci. Acad. Paris Ser. A 288, (1979), 237-239

[5] M.-C. Chang, A Filtered Bertini-type Theorem, J. reine und ang. Math. **397.** (1989), 214-219

[6] L. Chiodera and Ph. Ellia, Rank 2 Globally Generated Vector Bundles with $c_1 \leq 5$, arXiv:1111.5725v1 [mathAG]

[7] A. Grothendieck, Éléments de géométrie algébrique (EGA) I-III, IHES Publ. Math., Paris 1960, 1961, 1963

[8] E. Guardo and B. Harbourne: Resolutions of ideals of any six fat points in $\mathbb{P}^2$, J. Alg. **318**, 2, (2007), 619-640

[9] R. Hartshorne, Algebraic Geometry, Springer, 1977

[10] R. Hartshorne, Stable Vector Bundles of Rank 2 on $\mathbb{P}^3$, Math. Ann. **238,** (1978), 229-280

[11] G. Horrocks, Vector Bundles on the Punctured Spectrum of a Local Ring, Proc. London Math. Soc. **3**, 14, (1964), 689-713

[12] G. Horrocks, D. Mumford, A Rank 2 Vector Bundle of Rank 2 on $\mathbb{P}^4$ with 15000 Symmetries, Topology **12**, (1973), 63-81

[13] S. Huh, On Triple Embeddings of $\mathbb{P}^n$ in the Grassmannians, Math. Nachr. **284**, No 11-12, (2011), 1453-1461

[14] K. Hulek, Stable Rank-2 Vector Bundles on $\mathbb{P}_2$ with c_1 Odd, Math. Ann. **242**, (1979), 241-266

[15] P. Ionescu, Embedded Projective Varieties of Small Invariants, Proc. of the Week of Alg. Geom., Bucharest, LNM 1056, Springer, Berlin, Heidelberg, New-York, Tokyo (1984)

[16] N. M. Kumar, C. Peterson and A. P. Rao, Standard Vector Bundle Deformations on $\mathbb{P}^n$, Contemporary Mathematics **322,** "Hilbert Schemes, Vector Bundles and their interplay with Representation Theory (Columbia MO 2002), (2003), 151-163

[17] N. Manolache, Cohen-Macaulay Nilpotent Structures, Rev. Roum. Pures Appl. **31,** (1986), 563-575

[18] N. Manolache, Globally Generated Vector Bundles on $\mathbb{P}^3$ with $c_1 = 3$, arXiv:1202.5988v1 [math.AG]

[19] Ch. Okonek, 3-Mannigfaltigkeiten im $\mathbb{P}^5$ und ihre zugehörigen stabilen Garben, Manuscripta Math., **38**, (1982), 175-199

[20] Ch. Okonek, Moduli reflexiver Garben und Flächen von kleinem Grad in $\mathbb{P}^4$, Math. Z. **184**, (1983), 549-572

[21] Ch. Okonek, Über 2-codimensionale Untermannigfaltigkeiten vom Grad 7 in $\mathbb{P}^4$ und $\mathbb{P}^5$, Math. Z. **187**, (1984), 209-219

[22] Ch. Okonek, M. Schneider, H. Spindler, Vector Bundles on Complex Projective Spaces, in: Progress in Mathematics, vol. 3, Birkhäuser, Boston, Mass. 1980

[23] G. Ottaviani, Varietà proiettive di codimensione piccola, Quaderni INDAM, Aracne, Roma, 1995
[24] C. Peskine, L. Szpiro, Liaison des varietes algebriques, Inv. Math., **26.**, (1974), 271-302
[25] J.-P. Serre, Faisceaux algébriques cohérents, Ann. of Math. **61,** (1955), 197-278
[26] T. Sauer, Nonstable Reflexive Sheaves on $\mathbb{P}^3$, Trans. A.M.S. **281.** (1984), 633-655
[27] J. C. Sierra, A Degree Bound for Globally Generated Vector Bundles, Math. Z. **262,** no 3, (2009), 517-525
[28] J. C. Sierra, L. Ugaglia, On globally generated vector bundles on projective spaces, Journal of Pure and Applied Algebra **213,** (2009) 2141-2148.
[29] Swinnerton-Dyer, H.P.F., An enumeration of all varieties of degree 4, Amer. J. Math., **95**, (1973), 403-418
[30] H. Tango, An Example of Indecomposable Vector Bundle of Rank $n-1$ on $\mathbb{P}^n$, J. of Math. of Kyoto Univ., **16,** no. 1 (1976), 137-141
[31] G. Trautmann, Darstellung von Vektorraumbündeln über $\mathbb{C}^n - 0$, Archiv der Math. **24,** (1973), 303-313
[32] U. Vetter, Zu einem Satz von G. Trautmann über den Rang gewisser kohärenter analitischer Moduln, Archiv der Math. **24,** (1973), 158-161
[33] J. A. Vogelaar, Constructing Vector Bundles from Codimension-Two Subvarieties, PhD Thesis, Leiden, 1978
[34] * * * , Correspondence, Amer. J. Math., **79**, (1957), 951-952 (A. Weil, Ouvres Sc., II, 555-556)

Institute of Mathematics of the Romanian Academy *Simion Stoilow* P.O. Box 1-764, 014700 Bucharest, Romania